

Fig. 1





Fig. 2



STEP 1: SET UP 3 PCR REACTIONS:


PRODUCTS:

TUBE 1:

TUBE 2:

TUBE 3:


STEP 2: SET UP PCR REACTION WITH PRODUCTS OF TUBE 1 +  
PRODUCTS TUBE 2 + F1 + R4.




STEP 3: REPEAT STEP 2 USING PRODUCT FROM STEP 2 + PRODUCT  
FROM STEP 1, TUBE 3 + PRIMERS F1 + R3.

Fig. 3


6



STEP 1: SET UP 3 PCR REACTIONS:


TUBE 1: 

TUBE 2: 

TUBE 3: 

STEP 2: DIGEST PRODUCTS FROM STEP 1 WITH SUITABLE RESTRICTION ENDONUCLEASES.

STEP 3: LIGATE DIGESTED PRODUCT FROM STEP 2, TUBE 2 WITH DIGESTED PRODUCT FROM STEP 2, TUBE 1.



STEP 4: AMPLIFY VIA PCR LIGATED PRODUCTS OF STEP 3 WITH F1 + R4.



STEP 5: DIGEST AMPLIFIED PRODUCT OF STEP 4 WITH RESTRICTION ENDONUCLEASE #2.



STEP 6: LIGATE PRODUCT FROM STEP 5 WITH PRODUCT FROM STEP 2, TUBE 3.



STEP 7: AMPLIFY PRODUCT FROM STEP 6 WITH F1 + R3.

Fig. 4




Fig. 5